





GIG  
CYMRU  
NHS  
WALES

# FLU

**FLU IS A KILLER**

**Flu vaccination 2015-16**





## Why is flu serious?

Flu spreads easily and can cause serious illnesses which need to be treated in hospital. There are outbreaks every year, usually in winter. Flu is not the same as a bad cold. It causes many deaths.

## Does the vaccine work?

Yes, the vaccine usually prevents flu in 5 to 7 people in every 10 who have it. But as flu viruses change, the vaccine is changed every winter to match the new flu viruses.

## How will I know if I have flu?

When people get flu they tend to become ill quite quickly. Symptoms include fever, chills, headaches and aching muscles, often with a cough and sore throat. Often people who get flu feel so ill they need to stay in bed for a few days.

## What should I do if I get flu?

Use a tissue when you sneeze or cough and wash your hands or use hand sanitizer to help stop flu spreading.

Antibiotics won't help.

There are medicines called antivirals that can be given to those most at risk from flu. These may help if taken within 2 days of the start of the illness. If you are at increased risk of flu and its complications, and think you may have it, you should talk to your doctor urgently.

If you are in a risk group and haven't had the vaccine this flu season, you should have it as soon as you are feeling better. This will protect you against other types of flu.

## **What harm can flu do?**

Flu can lead to serious illnesses such as bronchitis and pneumonia that may need hospital treatment. Flu is more serious if you have certain long term illnesses, are pregnant or are 65 or older. In a normal winter, thousands of people will die from flu related illnesses in the UK.

## **Am I at greater risk from the effects of flu?**

Even if you feel healthy, if you have one of the following conditions you are at increased risk of complications from flu and should have a flu vaccination:

- A heart problem.
- A chest complaint or breathing difficulties, including asthma that requires regular steroid inhalers or tablets.
- Kidney disease.

- Lowered immunity due to disease or treatment (such as steroid medication or cancer treatment).
- Liver disease.
- Had a stroke or a transient ischaemic attack (mini-stroke).
- Diabetes.
- A neurological condition, for example multiple sclerosis (MS), cerebral palsy or post-polio syndrome.
- A problem with your spleen, for example sickle cell disease, or you have had your spleen removed.

**You should also have the flu vaccination if you are:**

- Pregnant.
- Aged 65 years or over.
- Living in a residential or nursing home.

## **Carers**

If you care for someone whose health would be at risk if you became ill with flu, you should have a free flu vaccine. This will help protect you and the person you care for.

You should also have the flu vaccine if you are a volunteer carer, a member of an organisation providing planned first aid or a community first responder.

## Health and social care workers

If you are a health or social care worker with face to face contact with patients or clients, you should have a flu vaccination to protect yourself and the people you care for. Your employer is responsible for paying for your vaccination.

## Pregnant women

If you are pregnant you should have the flu vaccine to protect yourself and your unborn child. If a pregnant woman gets flu, her baby is more likely to be born early or with a low birth weight, be stillborn or die within their first week. The vaccine also protects the baby in the first 4-6 months of life when flu can be very serious.

## Children

Children from 6 months of age with any of the conditions listed above or who have previously been admitted to hospital with a chest infection, should have a flu vaccination.

A nasal spray vaccine is being introduced for children from 2 to 16 years of age. It is more effective than the flu injection in children. This will help prevent children from catching flu and also prevent them from spreading it to others. There is a separate children's flu leaflet which provides the most recent information.

## **I had a flu vaccine last year. Do I need one this year?**

Yes, if you are in a risk group for flu you need to have a flu vaccine every year.

## **Is there anyone who should not have the flu vaccine?**

There are very few people who cannot have a flu vaccine. People who have previously had a serious allergic reaction to a flu vaccine (or any part of it) should not have that flu vaccine again.

Tell the nurse or doctor if you have a serious egg allergy. You can still be vaccinated but they need to know because some flu vaccines are not suitable.

The nasal spray vaccine should not be given to anyone who is under 2 or over 17 years, has a weakened immune system, severe asthma, a wheezy chest when the vaccine is due, or has previously received intensive care for an allergic reaction to eggs.

People who are unwell with a high temperature should delay vaccination until they are feeling better.

A runny nose, cough or other minor illness are not reasons to delay the vaccine.

## **Will I get any side effects?**

The flu vaccine is very safe. After having the vaccine, some people get a slight temperature, feel tired, have a headache, or aching muscles for a day or two.

After the injection, your arm may be a bit red and sore but other reactions are less common.

The nasal spray may cause a runny or blocked nose.

Other side effects are very rare.

Flu vaccines are quick and safe and could prevent weeks of serious illness.

## **Will the flu vaccine give me flu?**

No, the flu vaccine will not give you flu.

## **Will the flu vaccine protect me completely?**

The vaccine usually offers good protection against flu in the flu season it is given. Protection starts around two weeks after the vaccination.

You may still get flu even after vaccination but often with milder symptoms. Flu vaccines do not protect against colds and other flu-like illnesses.

## **What do I need to do next?**

If you are in any of the groups recommended to have a flu vaccine, it is important that you have it. Adults can have the vaccine free at their GP practice or at some pharmacies. If your child is in the age range for routine vaccination, you should be contacted by your GP or school nursing service. You can get advice from your GP practice, local pharmacy or NHS Direct Wales.

## Where can I get more information?

You can find out more about the vaccine, including vaccine contents and potential side effects, at [www.medicines.org.uk/emc/](http://www.medicines.org.uk/emc/) by entering the name of the vaccine in the search box and looking at the patient information leaflet (PIL).

The routine immunisation schedule that shows you which immunisations are routinely offered in Wales is available from [www.publichealthwales.org/routineimmunisation-schedule-wales](http://www.publichealthwales.org/routineimmunisation-schedule-wales)

If you have any questions or want more information, talk to your doctor or nurse, call NHS Direct Wales on 0845 4647 or visit

[www.nhsdirect.wales.nhs.uk/](http://www.nhsdirect.wales.nhs.uk/)



To order more copies of this leaflet, email: [hplibrary@wales.nhs.uk](mailto:hplibrary@wales.nhs.uk) or telephone 08456 064050.

**This leaflet is produced by:**

Public Health Division, Welsh Government  
and Vaccine Preventable Disease Programme  
Public Health Wales

[www.beatflu.org](http://www.beatflu.org)

